Orencia® Referral Form

PHONE: 1-800-277-7302 FAX: 1-866-374-6663 Email: intake@healthcoga.com ☐ ***Initial Order*** ☐ ***Reorder***

Demographics Information: Today's Date _____

Patient Name: _____ DOB: _____

Address: _____ City: _____ State: GA Zip: _____

Phone #: _____ ☐ Cell Alternate Phone #: _____ ☐ Cell

Height:_____ in/ft Weight: _____ lbs/kg Date weight recorded: _____ Last 4 of SSN: _____

Allergies: _____ ☐ **NKDA**

Diagnosis Information: ☐ Rheumatoid Arthritis M05.79

☐ Other Diagnosis/ICD10Code_____

Date Infusion to Begin: _____

Previous Infusion Reaction: ☐ YES ☐ NO

List Reactions: _____

Is this a first infusion? ☐ YES ☐ NO

Date of Last Infusion: _____

List of Failed Therapies: _____

Assessment Questions: *Please Provide Appropriate Documentation*

Diabetes? ☐ Yes ☐ No HTN? ☐ Yes ☐ No Cardiac History? ☐ Yes ☐ No Difficulty Breathing? ☐ Yes ☐ No
Requires Oxygen? ☐ Yes ☐ No Confusion /Disorientation? ☐ Yes ☐ No Neuro History? ☐ Yes ☐ No
Date of Negative TB test: _____

ORENCIA (abatacept): J0129 Patient Dose per Body Weight

☐ < 60 kg: 500 mg IV over 30 minutes at 0, 2, and 4 weeks: then every 4 weeks for _____ months
☐ 60 to 100 kg: 750 mg IV over 30 minutes at 0, 2, and 4 weeks: then every 4 weeks for _____ months.
☐ >100 kg: 1000 mg IV over 30 minutes at 0, 2, and 4 weeks: then every 4 weeks for _____ months.

Pre-Medications: Patient to Provide and take 30 minutes prior to infusion. *Will receive 50ml NS pre- and post- infusion.*

☐ Benadryl 25 MG PO x1 ☐ Benadryl 50 mg PO x1
☐ Acetaminophen 325 mg PO x1 ☐ Acetaminophen 650 mg PO x1
☐ Other Pre-medication: _____ Follow IV Care Anaphylaxis Protocol

Access: ☐**PIV** ☐**PORT** ☐ **PICC** **Route:** ☐ **IV**

Catheter Care/Flush: To gain access, use for any ordered pre-medications, and/or PRN to maintain access and patency

PIV – NS 5ml: Qty 3

PORT – NS 10ml: Qty 3, Heparin 5mL 100units/mL x 1 per infusion.

PICC – NS 10ml: Qty 3, Heparin 5mL 10units/mL x 1 per infusion

Supplies: (please strike through if not required)

Administration Supplies (A4222) – 1 per infusion
Catheter Care Supplies (A4221) – 1 per week
Infusion Pump (E0781)
Nursing services to administer medications

Administration through an in-line 0.2- to 1.2-micron, sterile, nonpyrogenic, low-protein binding filter is required.

Administer the anaphylaxis protocol for adverse reactions

Provider signature: _____

NPI # _____ Phone _____ Fax _____

Office Address: _____ City: _____ St ____Zip: _____

Printed MD Name: _____ Contact Name: _____

☐ Demographic information ☐ Patient Ins. Card ☐ H & P ☐ Office notes ☐ Failed therapies in documentation
☐ Up to date medication list

9/23 TSP